| レジメンNO 抗TNFα抗体(初回 ~3コース) | | | | | | | | | | | | | レジメン名 | | | TNF | α抗体 | ▶(初回 | □~ 3: | 3 —7 | ス) | | | | | | | | | | | | | | | | | | | | |
|--------------------------|--------|--------------------|------|-------|-------|-----|--|-----|-----|---|---|---|-------|---|----|-----|------|--------|--------------|-------------|------|-----|------|----|-----|----|------|-----|-----|-------------|----|----|------|------|---------------|----|------|------|------|----|--|
| | 癌和 | 遺瘍性大陽炎&ク ローン病 | | | | | | | | | | | | | 診療 | 春科 | þ | 内科 | | | | | | =- | -ス日 | 数 | 98日 | | | ⊐ –; | ス数 | ŧ | 栗準[- |] # | k 小[-] | 最: | 大[-] | | | | |
| | 77 D.I | | W. E | | 8877 | | | 1W | | | | | 2W | | | | | 3W | | | | | | 4 | | 4W | | | | 5 | 5W | | | 6W | | | | | | | |
| 10 | 種別 | 内容 | 数量 | 単位 | 開始 | 時間 | | 1 2 | 2 3 | 4 | 5 | 6 | 7 8 | 9 | 10 | 11 | 12 1 | 3 14 | 15 | 16 | 17 1 | 8 1 | 9 20 | 21 | 22 | 23 | 24 2 | 5 2 | 6 2 | 7 28 | 29 | 30 | 31 | 32 3 | 33 34 | 35 | 36 | 37 3 | 8 39 | 40 | |
| Ť | | 生理食塩液「ヒカリ」 50mL | | 瓶 | | | | | | | | | | | | | Ť | | | | | | | | | | | Ť | Ť | | | | | Ŧ | | П | П | T | T | - | |
| 1 | 注射 | ブライミング | | | 00:00 | - | | Э | | | | | | | | | | | 0 | | | | | | | | | | | | | | | | | | | | | | |
| | | 静脈内注射 | | 1 🗇 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | 2時間以上で | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | 注射 | 生理食塩液「ヒカリ」 250mL | | l 袋 | 00:00 | .00 | | 0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 | 注别 | (癌)インフリキシマブBS点滴静注用 | : | mg/Kg | 00:00 | - | | ار | | | | | | | | | | | | 0 | | | | | | | | | | | | | | | | | | 1 | | | |
| | | 静脈内注射 | | 1 🗇 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | | i | | | | | |